

# Phospho-FAK-Y925 Rabbit pAb

货号: **B14985** 

#### 产品信息

反应 Human	
宿主 Rabbit	
克隆性 Polyclonal	
预测反应 WB: Human fibrosarcoma cells	5
应用 WB	
推荐浓度 WB: 1:500 - 1:2000	
理论分子量 39kDa/48kDa/63kDa/99kDa/114	4kDa/119kDa/120kDa
实测分子量 125kDa	
形式 Liquid	
保存条件 Store at -20°C. Avoid freeze / th Buffer: PBS with 0.02% sodium	
偶联物 Unconjugated	
阳性对照 JK	
	toplasm,Cytoplasmic side,Nucleus,Peripheral membrane protein,cell corte al adhesion,microtubule organizing center
纯化 Affinity purification	

### 抗原信息

抗原信息	A synthetic phosphorylated peptide around Y925 of human FAK (NP_722560.1).
序列	KVYEN


#### 靶点信息

劬	Ŧ	ù	育	京

This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesio ns that form between cells growing in the presence of extracellular matrix constituents. The encoded prot ein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Activation of this gene may be an important early step in cell growth a nd intracellular signal transduction pathways triggered in response to certain neural peptides or to cell int eractions with the extracellular matrix. Several transcript variants encoding different isoforms have been found for this gene.

基因ID	5747
基因名	PTK2
Swiss	Q05397
别名	FADK;FAK1;FRNK;PPP1R71;p125FAK;pp125FAK;PTK2

## 产品验证



Western blot analysis of Phospho-FAK-Y925 expressed in JK using Phospho-FAK-Y925 Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn