

Phospho-ZAP70-Y319 Rabbit pAb

货号: B15062

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:50 - 1:100
理论分子量	35kDa/55kDa/69kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,Preincubated with blocking peptide
细胞定位	Cell membrane,Cytoplasm,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	A synthetic phosphorylated peptide around Y319 of human ZAP70 (NP_001070.2).
序列	SPYSD

靶点信息

This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upo n T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transducti on in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte d evelopment. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency di sease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode di fferent isoforms have been found for this gene.

基因ID	7535
基因名	ZAP70
Swiss	P43403
别名	ZAP70;ADMIO2;IMD48;SRK;STCD;STD;TZK;ZAP-70

产品验证



Western blot analysis of Phospho-ZAP70-Y319 expressed in Jurkat, Preincubated with blocking peptide us ing Phospho-ZAP70-Y319 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL En hanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn