

MonoMethyl-H2B-E105 Rabbit pAb

货号: B15066

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:100 - 1:500
理论分子量	14kDa
实测分子量	15KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,NIH/3T3
细胞定位	cytosol,nucleoplasm,nucleus
纯化	Affinity purification

抗原信息

抗原信息	A synthetic monomethylated peptide around E105 of human H2B (NP_066406.1).
序列	LPGEL


靶点信息

研究背景

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octa mer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is fu rther compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes t o form higher order chromatin structures. This gene is intronless and encodes a replication-dependent his tone that is a member of the histone H2B family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chrom osome 6p22-p21.3.

基因ID	3018
基因名	HIST1H2BB
Swiss	P33778
别名	H2B.1; H2B/f; H2BFF; HIST1H2BB

产品验证



Western blot analysis of MonoMethyl-H2B-E105 expressed in HeLa,NIH/3T3 using MonoMethyl-H2B-E 105 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/ proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn