

# SARS-CoV-2 NSP13 Rabbit pAb

货号: B15068

## 产品信息

反应	SARS-CoV-2
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:2000 - 1:6000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	
实测分子量	75KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-601 of coronavirus NS P13 (YP_009725308.1).
序列	AVGACVL CNSQ TSLRCGACIRPFL CCKCCYDH VISTSHKL VL SVNP YVCNAP GCDV TDVT QLYL GGMSYY CKSH KPPIS FPLC ANGQV FGLY KNTCV GSDN VTDF NAIAT CDWT NAGD YILANT CTER LKLF AAET LKATE ETFK LSYG IATV REVLS DREL HLSWEVGKPRP PLNR NYVFTGY RVT KNS KVQIGE YT FEKGDY GDAV VYRGTT TYKLN VGDY FVLT SHTV MPLS APTL VPQ EHYVR ITGLY PTLN ISDEF SSNV ANYQ KVGM QKYSTL QGPPGTGKSHFAIGL ALY YPSARIV YTACCS HAAVD ALCE KALKY LP IDKCSRII PARAR VECFD KFKVN STLE QYVF CTVN ALPET TADIVV FDEISM ATNY DL SVVN AR LRAK HYVY IGDP AQLP AP RT LLTK GTLE PEYF NSV CRLM KTI GPDM FLGT CRRCP AEIV DT VSAL VYDN KLKAH KDKSA QCFC KMFY KGVI THDV SAIN RPQ IGVVREFL TRNP AWRK AVFISP YNSQ NAVASK IGLPLT QTV DSSQ GSEY DYVIFT QTTE AHSC NVRN VAITRA KV GILCI MSDR DLYD KQFT SLEIP RR NVATLQ

## 靶点信息

研究背景	[Spike protein S1]: attaches the virion to the cell membrane by interacting with host receptor, initiating the infection. Binding to human ACE2 receptor and internalization of the virus into the endosomes of the host cell induces conformational changes in the Spike glycoprotein. Binding to host NRP1 and NRP2 via C-terminal polybasic sequence enhances virion entry into host cell. This interaction may explain virus tropism of human olfactory epithelium cells, which express high level of NRP1 and NRP2 but low level of ACE2. The stalk domain of S contains three hinges, giving the head unexpected orientational freedom. Uses human TMPRSS2 for priming in human lung cells which is an essential step for viral entry. Can be alternatively processed by host furin. Proteolysis by cathepsin CTSL may unmask the fusion peptide of S2 and activate membranes fusion within endosomes.
基因ID	43740578
基因名	
Swiss	P0DTD1
别名	

## 产品验证



Western blot analysis of SARS-CoV-2 NSP13 expressed in 293T using SARS-CoV-2 NSP13 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)