

Phospho-MARCKS-S158 Rabbit pAb

货号: B15073

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:100 - 1:200
理论分子量	31kDa
实测分子量	85kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3
细胞定位	Cytoplasm,Lipid-anchor,Membrane,cytoskeleton
纯化	Affinity purification

抗原信息


抗原信息	A phospho specific peptide corresponding to residues surrounding S158 of human MARCKS
序列	

靶点信息

研究背景	The protein encoded by this gene is a substrate for protein kinase C. It is localized to the plasma membrane and is an actin filament crosslinking protein. Phosphorylation by protein kinase C or binding to calcium-calmodulin inhibits its association with actin and with the plasma membrane, leading to its presence in the cytoplasm. The protein is thought to be involved in cell motility, phagocytosis, membrane trafficking and mitogenesis.
基因ID	4082

基因名	MARCKS
Swiss	P29966
别名	MARCKS;80K-L;MACS;PKCSL;PRKCSL

产品验证



Western blot analysis of Phospho-MARCKS-S158 expressed in NIH/3T3 using Phospho-MARCKS-S158 R abbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure ti me: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn