

Phospho-CDKN1B/p27KIP1-T187 Rabbit pAb

货号: B15096

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:100
理论分子量	22kDa
实测分子量	26kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cytoplasm,Endosome,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	A phospho specific peptide corresponding to residues surrounding T187 of human CDKN1B/p27KIP1
序列	

靶点信息

研究背景	This gene encodes a cyclin-dependent kinase inhibitor, which shares a limited similarity with CDK inhibitor CDKN1A/p21. The encoded protein binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state. Mutations in this gene are associated with multiple endocrine neoplasia type IV (MEN4).
基因ID	1027

基因名	CDKN1B
Swiss	P46527
别名	CDKN1B;CDKN4;KIP1;MEN1B;MEN4;P27KIP1;p27 KIP 1

产品验证



Western blot analysis of Phospho-CDKN1B/p27KIP1-T187 expressed in HeLa using Phospho-CDKN1 B/p27KIP1-T187 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5 000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn