

# **TPSAB1** Rabbit pAb

货号**: B15142** 

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	29kDa/30kDa
实测分子量	30KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse skin,Rat skin
细胞定位	Secreted
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human TPSAB1 (NP_003285 .2).	
序列	MLNLLLLALPVLASRAYAAPAPGQALQRVGIVGGQEAPRSKWPWQVSLRVHGPYWMHFCGGSLIHPQWVLTAAHCVGP DVKDLAALRVQLREQHLYYQDQ	

靶点信息

研究背景	Tryptases comprise a family of trypsin-like serine proteases, the peptidase family S1. Tryptases are enzy matically active only as heparin-stabilized tetramers, and they are resistant to all known endogenous prot einase inhibitors. Several tryptase genes are clustered on chromosome 16p13.3. These genes are charact erized by several distinct features. They have a highly conserved 3' UTR and contain tandem repeat sequ ences at the 5' flank and 3' UTR which are thought to play a role in regulation of the mRNA stability. Thes e genes have an intron immediately upstream of the initiator Met codon, which separates the site of trans cription initiation from protein coding sequence. This feature is characteristic of tryptases but is unusual in other genes. The alleles of this gene exhibit an unusual amount of sequence variation, such that the all eles were once thought to represent two separate genes, alpha and beta 1. Beta tryptases appear to be the main isoenzymes expressed in mast cells; whereas in basophils, alpha tryptases predominate. Tryptases have been implicated as mediators in the pathogenesis of asthma and other allergic and inflammatory disorders.
基因ID	7177
基因名	TPSAB1
Swiss	Q15661
别名	TPSAB1;TPS1;TPS2;TPSB1

## 产品验证



Western blot analysis of TPSAB1 expressed in Mouse skin,Rat skin using TPSAB1 Rabbit pAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking b uffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn