

APOBEC3F Rabbit pAb

货号**: B15150**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	9kDa/11kDa/45kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,BxPC-3,HepG2
细胞定位	Cytoplasm,P-body
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-65 of human APOBEC 3F (NP_660341.2).
序列	MKPHFRNTVERMYRDTFSYNFYNRPILSRRNTVWLCYEVKTKGPSRPRLDAKIFRGQVYSQPEHH

靶点信息

	研究背景	This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudo genes found in a cluster, thought to result from gene duplication, on chromosome 22. Members of the clu ster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deam inase APOBEC1. It is thought that the proteins may be RNA editing enzymes and have roles in growth or c ell cycle control. Alternatively spliced transcript variants encoding different isoforms have been identified .
--	------	---

基因ID	200316
基因名	APOBEC3F
Swiss	Q8IUX4
别名	APOBEC3F;A3F;ARP8;BK150C2.4.MRNA;KA6

产品验证



Western blot analysis of APOBEC3F expressed in HeLa,BxPC-3,HepG2 using APOBEC3F Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per l ane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn