

H6PD Rabbit pAb

货号: **B15234**

产品信息

| 反应 | Human,Mouse,Rat |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 88kDa |
| 实测分子量 | 89kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Mouse liver, Mouse kidney |
| 细胞定位 | Endoplasmic reticulum lumen |
| 纯化 | Affinity purification |

抗原信息


| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 20-200 of human H6PD (NP_004276.2). | |
|------|---|--|
| 序列 | QELQGHVSIILLGATGDLAKKYLWQGLFQLYLDEAGRGHSFSFHGAALTAPKQGQELMAKALESLSCPKDMAPSHCAEH KDQFLQLSQYRQLKTAEDYQALNKDIEAQLQHAGLREAGRIFYFSVPPFAYEDIARNINSSCRPGPGAWLRVVLEKPFGHD HFSAQQLATELGTFFQEEEMY | |

靶点信息

| 研究背景 | There are 2 forms of glucose-6-phosphate dehydrogenase. G form is X-linked and H form, encoded by this gene, is autosomally linked. This H form shows activity with other hexose-6-phosphates, especially galact ose-6-phosphate, whereas the G form is specific for glucose-6-phosphate. Both forms are present in most tissues, but H form is not found in red cells. |
|--------------|---|
| 基因 ID | 9563 |

| 基因名 | H6PD |
|-------|------------------------|
| Swiss | O95479 |
| 别名 | H6PD;CORTRD1;G6PDH;GDH |

产品验证



Western blot analysis of H6PD expressed in Mouse liver, Mouse kidney using H6PD Rabbit pAb at 1:1000. S econdary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn