

ACTC1 Rabbit pAb

货号: **B15236**

产品信息

Human,Mouse,Rat
Rabbit
Polyclonal
WB IHC
WB: 1:500 - 1:1000 IHC: 1:50 - 1:200
42kDa
42KDa
Liquid
Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
Unconjugated
C2C12,H9C2,RD,Mouse lung,Mouse heart,Rat lung,Rat heart
Cytoplasm,cytoskeleton
Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human ACTC1 (NP_005150. 1).
序列	MCDDEETTALVCDNGSGLVKAGFAGDDAPRAVFPSIVGRPRHQGVMVGMGQKDSYVGDEAQSKRGILTLKYPIEHGIIT NWDDMEKIWHHTFYNELRVAP

靶点信息

研究背景	Actins are highly conserved proteins that are involved in various types of cell motility. Polymerization of gl obular actin (G-actin) leads to a structural filament (F-actin) in the form of a two-stranded helix. Each actin can bind to four others. The protein encoded by this gene belongs to the actin family which is comprise d of three main groups of actin isoforms, alpha, beta, and gamma. The alpha actins are found in muscle ti ssues and are a major constituent of the contractile apparatus. Defects in this gene have been associated with idiopathic dilated cardiomyopathy (IDC) and familial hypertrophic cardiomyopathy (FHC).
------	---

基因ID	70
基因名	ACTC1
Swiss	P68032
别名	ACTC1;ACTC;ASD5;CMD1R;CMH11;LVNC4;actin

产品验证



Western blot analysis of ACTC1 expressed in Mouse lung,Rat lung,C2C12 using ACTC1 Rabbit pAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn