

SRGN Rabbit pAb

货号: **B15262**

产品信息

反应 Hum	nan
	IMIT
宿主 Rabb	bit
克隆性 Polyo	rclonal
预测反应	
应用 WB	IHC
	: 1:500 - 1:2000 : 1:50 - 1:200
理论分子量 17kC	Da
实测分子量 17kC	Da
形式 Liqui	iid
	re at -20°C. Avoid freeze / thaw cycles. Fer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物 Uncc	onjugated
阳性对照 THP-	-1
细胞定位 Cyto	pplasmic granule,Golgi apparatus,Secreted,extracellular space
纯化 Affin	nity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 28-158 of human SRGN (NP_002718.2).
序列	YPTRRARYQWVRCNPDSNSANCLEEKGPMFELLPGESNKIPRLRTDLFPKTRIQDLNRIFPLSEDYSGSGFGSGSGSGSGSGSGSGFLTEMEQDYQLVDESDAFHDNLRSLDRNLPSDSQDLGQHGLEEDFML

靶点信息

研究背景	This gene encodes a protein best known as a hematopoietic cell granule proteoglycan. Proteoglycans stor ed in the secretory granules of many hematopoietic cells also contain a protease-resistant peptide core, which may be important for neutralizing hydrolytic enzymes. This encoded protein was found to be associ ated with the macromolecular complex of granzymes and perforin, which may serve as a mediator of granule-mediated apoptosis. Two transcript variants, only one of them protein-coding, have been found for this gene.
------	--

基因ID	5552
基因名	SRGN
Swiss	P10124
别名	SRGN;PPG;PRG1;serglycin

产品验证



Western blot analysis of SRGN expressed in THP-1 using SRGN Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dr y milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn