

BTK Rabbit pAb

货号: B15380

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100
理论分子量	76kDa/79kDa
实测分子量	76kDa/79kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Daudi cells, mouse spleen tissue, L02 cells, Ramos cells
细胞定位	Cell membrane,Cytoplasm,Nucleus,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 30-240 of human BTK (NP_000052.1).
序列	FLLTVHKLSYYEYDFERGRRGSKKGSIDVEKITCVETVVPEKNPPPERQIPRRGEESSEMEQISIIERFPYPFQVVYDEGPLYV FSPTEELRKRWIHLKKNVIRYNSDLVQKYHPCFWIDGQYLCCSQTAKNAMGCQILENRNGSLKPGSSHRKTKKPLPPTPEE DQILKKPLPPEPAAAPVSTSELKKVVALYDYMPMNANDLQLRKGDE

靶点信息

研究背景	The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement. Alternative splicing results in multiple transcript variants encoding different isoforms.
------	---

基因ID	695
基因名	BTK
Swiss	Q06187
别名	AGMX1;AT;ATK;BPK;IMD1;PSCTK1;XLA;BTK

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn