

SKAP2 Rabbit pAb

货号: **B15381**

产品信息

反应	Human, Mouse, Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	41kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-120 of human SKAP2 (NP_003921.2).
序列	MPNPSSTSSPYPLPEEIRNLLADVETFVADILKGENLSKKAKEKRESLIKKIKDVKSIYLQEFQDKGDAEDGEEYDDPFAGPP DTISLASERYDKDDEAPSDGAQFPPIAAQDLPFVLKA

靶点信息

研究背景	The protein encoded by this gene shares homology with Src kinase-associated phosphoprotein 1, and is a substrate of Src family kinases. It is an adaptor protein that is thought to play an essential role in the Src signaling pathway, and in regulating proper activation of the immune system. This protein contains an a mino terminal coiled-coil domain for self-dimerization, a plecskstrin homology (PH) domain required for in teractions with lipids at the membrane, and a Src homology (SH3) domain at the carboxy terminus. Some reports indicate that this protein inhibits actin polymerization through interactions with actin assembly fa ctors, and might negatively regulate the invasiveness of tumors by modulating actin assembly. Alternative splicing results in multiple transcript variants encoding different isoforms.
基因 ID	8935
基因名	SKAP2
Swiss	O75563
别名	SKAP2;PRAP;RA70;SAPS;SCAP2;SKAP-HOM;SKAP55R

产品验证



Western blot analysis of SKAP2 expressed in Mouse lung using SKAP2 Rabbit pAb at 1:1000. Secondary an tibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% n on-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn