

PTPRK Rabbit pAb

货号: **B15466**

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	162kDa/163kDa/164kDa
实测分子量	162kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver
细胞定位	Cell junction,Cell membrane,Single-pass type I membrane protein,adherens junction
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 600-750 of human PTPR K (NP_001129120.1).	
序列	EGVDASLNETATTITVLLRPAQAKGAPISAYQIVVEELHPHRTKREAGAMECYQVPVTYQNAMSGGAPYYFAAELPPGNLPE PAPFTVGDNRTYQGFWNPPLAPRKGYNIYFQAMSSVEKETKTQCVRIATKAAATEEPEVIPDPAKQTDR	

靶点信息

研究背景	The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP possesses an extracellular region, a single transmembrane region, and two tandem catalytic domains, and thus represents a receptor-type PT P. The extracellular region contains a meprin-A5 antigen-PTP mu (MAM) domain, an Ig-like domain and fo ur fibronectin type III-like repeats. This PTP was shown to mediate homophilic intercellular interaction, po ssibly through the interaction with beta- and gamma-catenin at adherens junctions. Expression of this ge ne was found to be stimulated by TGF-beta 1, which may be important for the inhibition of keratinocyte p roliferation.
基因 ID	5796
基因名	PTPRK
Swiss	Q15262
别名	PTPRK;R-PTP-kappa

产品验证



Western blot analysis of PTPRK expressed in Mouse liver using PTPRK Rabbit pAb at 1:1000. Secondary an tibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% n on-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn