

F13A1 Rabbit pAb

货号: **B15474**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	83kDa
实测分子量	83kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,SH-SY5Y,BxPC-3,Mouse testis,Rat brain,Rat testis
细胞定位	Cytoplasm,Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 600-732 of human F13A 1 (NP_000120.2).	
序列	LEQASLHFFVTARINETRDVLAKQKSTVLTIPEIIIKVRGTQVVGSDMTVTVEFTNPLKETLRNVWVHLDGPGVTRPMKKMF REIRPNSTVQWEEVCRPWVSGHRKLIASMSSDSLRHVYGELDVQIQRRPSM	

靶点信息

研究背景	This gene encodes the coagulation factor XIII A subunit. Coagulation factor XIII is the last zymogen to bec ome activated in the blood coagulation cascade. Plasma factor XIII is a heterotetramer composed of 2 A s ubunits and 2 B subunits. The A subunits have catalytic function, and the B subunits do not have enzymat ic activity and may serve as plasma carrier molecules. Platelet factor XIII is comprised only of 2 A subunit s, which are identical to those of plasma origin. Upon cleavage of the activation peptide by thrombin and i n the presence of calcium ion, the plasma factor XIII dissociates its B subunits and yields the same active enzyme, factor XIIIa, as platelet factor XIII. This enzyme acts as a transglutaminase to catalyze the forma tion of gamma-glutamyl-epsilon-lysine crosslinking between fibrin molecules, thus stabilizing the fibrin cl ot. It also crosslinks alpha-2-plasmin inhibitor, or fibronectin, to the alpha chains of fibrin. Factor XIII defici ency is classified into two categories: type I deficiency, characterized by the lack of both the A and B subu nits; and type II deficiency, characterized by the lack of the A subunit alone. These defects can result in a lifelong bleeding tendency, defective wound healing, and habitual abortion.
基因 ID	2162
基因名	F13A1
Swiss	P00488
别名	F13A1;F13A

产品验证



Western blot analysis of F13A1 expressed in Mouse testis,Rat brain,Jurkat using F13A1 Rabbit pAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn