

IFIT3 Rabbit pAb

货号: B15513

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	55kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,COS-7,Mouse liver
细胞定位	Cytoplasm, Mitochondrion
纯化	Affinity purification

抗原信息

抗原信息	A synthetic peptide of mouse IFIT3
序列	


靶点信息

研究背景

IFN-induced antiviral protein which acts as an inhibitor of cellular as well as viral processes, cell migration , proliferation, signaling, and viral replication. Enhances MAVS-mediated host antiviral responses by servi ng as an adapter bridging TBK1 to MAVS which leads to the activation of TBK1 and phosphorylation of IRF 3 and phosphorylated IRF3 translocates into nucleus to promote antiviral gene transcription. Exhibits an a ntiproliferative activity via the up-regulation of cell cycle negative regulators CDKN1A/p21 and CDKN1B/p 27. Normally, CDKN1B/p27 turnover is regulated by COPS5, which binds CDKN1B/p27 in the nucleus and exports it to the cytoplasm for ubiquitin-dependent degradation. IFIT3 sequesters COPS5 in the cytoplasm , thereby increasing nuclear CDKN1B/p27 protein levels. Upregulates CDKN1A/p21 by downregulating MY C, a repressor of CDKN1A/p21. Can negatively regulate the apoptotic effects of IFIT2.

基因 ID	15959
基因名	
Swiss	
别名	IFIT3;CIG-49;GARG-49;IFI60;IFIT4;IRG2;ISG60;P60;RIG-G;cig41

产品验证



Western blot analysis of IFIT3 expressed in HT-29,COS-7,Mouse liver using IFIT3 Rabbit pAb at 1:1000. S econdary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blockin g buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn