

# **GNA12 Rabbit pAb**

货号: **B15522** 

### 产品信息

反应	Human, Mouse, Rat
۸. بدن	
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	34kDa/37kDa/44kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse pancreas,Rat pancreas
细胞定位	Lipid-anchor,Membrane
纯化	Affinity purification


### 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 285-381 of human GNA 12 (NP_031379.2).
序列	LFFNVSIILFLNKMDLLVEKVKTVSIKKHFPDFRGDPHRLEDVQRYLVQCFDRKRRNRSKPLFHHFTTAIDTENVRFVFHAV KDTILQENLKDIMLQ

靶点信息

研究背景	Guanine nucleotide-binding proteins (G proteins are involved as modulators or transducers in various tran smembrane signaling systems. Activates effector molecule RhoA by binding and activating RhoGEFs (ARH GEF12/LARG. GNA12-dependent Rho signaling subsequently regulates transcription factor AP-1 (activatin g protein-1 (By similarity. GNA12-dependent Rho signaling also regulates protein phosphatese 2A activati on causing dephosphorylation of its target proteins. Promotes tumor cell invasion and metastasis by activating RhoA/ROCK signaling pathway and up-regulating proinflammatory cytokine production. Inhibits CDH 1-mediated cell adhesion in process independent from Rho activation. Together with NAPA promotes CD H5 localization to plasma membrane. May play a role in the control of cell migration through the TOR sign aling cascade.
基因 <b>ID</b>	2768
基因名	GNA12
Swiss	Q03113
别名	GNA12;NNX3;RMP;gep

### 产品验证



Western blot analysis of GNA12 expressed in Mouse pancreas,Rat pancreas using GNA12 Rabbit pAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn