

# MRPS31 Rabbit pAb

货号: **B15566** 

## 产品信息


反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	45kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,HeLa,HepG2,MCF7,A-431
细胞定位	Mitochondrion
纯化	Affinity purification

## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 66-395 of human MRPS 31 (NP_005821.2).
序列	ICSKKDKQSVRTEETSKETSESQDSEKENTKKDLLGIIKGMKVELSTVNVRTTKPPKRRPLKSLEATLGRLRRATEYAPKKRI EPLSPELVAAASAVADSLPFDKQTTKSELLSQLQQHEEESRAQRDAKRPKISFSNIISDMKVARSATARVRSRPELRIQFDE GYDNYPGQEKTDDLKKRKNIFTGKRLNIFDMMAVTKEAPETDTSPSLWDVEFAKQLATVNEQPLQNGFEELIQWTKEGKL WEFPINNEAGFDDDGSEFHEHIFLEKHLESFPKQGPIRHFMELVTCGLSKNPYLSVKQKVEHIEWFRNYFNEKKDILKESNI QFN

研究背景	Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a l arge 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribo somes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryo tic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the m itoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy rec ognition by sequence homology. The 28S subunit of the mammalian mitoribosome may play a crucial and characteristic role in translation initiation. This gene encodes a 28S subunit protein that has also been ass ociated with type 1 diabetes; however, its relationship to the etiology of this disease remains to be clarified. Pseudogenes corresponding to this gene have been found on chromosomes 3 and 13.
基因ID	10240
基因名	MRPS31
Swiss	Q92665
别名	MRPS31;IMOGN38;MRP-S31;S31mt

## 产品验证



Western blot analysis of MRPS31 expressed in Jurkat, HeLa, HepG2 using MRPS31 Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit  $\lg G$  (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blo cking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn