

# Phospho-FOXO4-S197 Rabbit pAb

货号: B15608

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:100 - 1:500 <b>IHC:</b> 1:50 - 1:100 <b>IF/ICC:</b> 1:100 - 1:200
理论分子量	47kDa/53kDa
实测分子量	65KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293F,Jurkat
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic phosphorylated peptide around S197 of human FOXO4 (NP_005929.2).
序列	AASMD

## 靶点信息

研究背景	This gene encodes a member of the O class of winged helix/forkhead transcription factor family. Proteins encoded by this class are regulated by factors involved in growth and differentiation indicating they play a role in these processes. A translocation involving this gene on chromosome X and the homolog of the Dr osophila trithorax gene, encoding a DNA binding protein, located on chromosome 11 is associated with le ukemia. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	4303

基因名	FOXO4
Swiss	P98177
别名	AFX;AFX1;MLLT7;FOXO4

产品验证



Western blot analysis of Phospho-FOXO4-S197 expressed in 293F,Jurkat using Phospho-FOXO4-S197 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/prot eins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exp osure time: 120s.

实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)