

PAK4 Rabbit pAb

货号: **B15610**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	47kDa/48kDa/54kDa/64kDa
实测分子量	70KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LNCaP,Mouse testis,Mouse brain
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 450-550 of human PAK4 (NP_005875 .1).
序列	HDGRVKLSDFGFCAQVSKEVPRRKSLVGTPYWMAPELISRLPYGPEVDIWSLGIMVIEMVDGEPPYFNEPPLKAMKMIRDN LPPRLKNLHKVSPSLKGFLD

靶点信息

研究背景	PAK proteins, a family of serine/threonine p21-activating kinases, include PAK1, PAK2, PAK3 and PAK4. PA K proteins are critical effectors that link Rho GTPases to cytoskeleton reorganization and nuclear signalin g. They serve as targets for the small GTP binding proteins Cdc42 and Rac and have been implicated in a wide range of biological activities. PAK4 interacts specifically with the GTP-bound form of Cdc42Hs and w eakly activates the JNK family of MAP kinases. PAK4 is a mediator of filopodia formation and may play a r ole in the reorganization of the actin cytoskeleton. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.
基因ID	10298
基因名	PAK4
Swiss	096013
别名	PAK4

产品验证



Western blot analysis of PAK4 expressed in LNCaP,Mouse testis,Mouse brain using PAK4 Rabbit pAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn