

# MELK Rabbit pAb

货号: B15620

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:100 - 1:200
理论分子量	52kDa/59kDa/66kDa/69kDa/70kDa/71kDa/74kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,SKOV3,HT-29,HeLa,B cells,Mouse thymus,Mouse spleen,Rat testis
细胞定位	Cell membrane,Peripheral membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 412-651 of human MELK (NP_055606.1).
序列	LCRTPANKLKNKENVYTPKSAVKNEEYFMFPEKTPVNKNQHKREILTTPNRYTTPSKARNQCLKETPIKIPVNSTGTDKLMT GVISPERRCRSVELDLNQAHMEETPKRKGAKEVFGSLERGLDKVITVLTRSKRKGSARDGPRRLKLHYNVTTRLVNPDQLL NEIMSILPKKHVDFVQKGYTLKCQTQSDFGKVTMQFELEVQQLQKPDVVGIRRQRLKGDWVYKRLVEDILSSCKV

靶点信息

研究背景	Serine/threonine-protein kinase involved in various processes such as cell cycle regulation, self-renewal of stem cells, apoptosis and splicing regulation. Has a broad substrate specificity; phosphorylates BCL2L14 , CDC25B, MAP3K5/ASK1 and ZNF622. Acts as an activator of apoptosis by phosphorylating and activating MAP3K5/ASK1. Acts as a regulator of cell cycle, notably by mediating phosphorylation of CDC25B, promoting localization of CDC25B to the centrosome and the spindle poles during mitosis. Plays a key role in cell proliferation and carcinogenesis. Required for proliferation of embryonic and postnatal multipotent neural progenitors. Phosphorylates and inhibits BCL2L14, possibly leading to affect mammary carcinogenesis by mediating inhibition of the pro-apoptotic function of BCL2L14. Also involved in the inhibition of spliceosome assembly during mitosis by phosphorylating ZNF622, thereby contributing to its redirection to the nucleus. May also play a role in primitive hematopoiesis.
基因ID	9833
基因名	MELK
Swiss	Q14680
别名	MELK;HPK38

产品验证



实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)