

LDB2 Rabbit pAb

货号**: B15636**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	38kDa/39kDa/42kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Raji,Mouse kidney
细胞定位	Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 204-373 of human LDB2 (NP_001281.1).	
序列	LRLCVILEPMQELMSRHKTYNLSPRDCLKTCLFQKWQRMVAPPAEPTRQPTTKRRKRKNSTSSTSNSSAGNNANSTGSKK KTTAANLSLSSQVPDVMVVGEPTLMGGEFGDEDERLITRLENTQYDAANGMDDEEDFNNSPALGNNSPWNSKPPATQE TKSENPPPQASQ	

靶点信息

研究背景	The protein encoded by this gene belongs to the LIM-domain binding family. Members of this family are c haracterized by a conserved nuclear localization sequence, an amino-terminal homodimerization domain and a carboxy-terminal LIM interaction domain. These proteins function as adapter molecules to allow as sembly of transcriptional regulatory complexes. Genetic association studies suggest functions for this gen e in rhegmatogenous retinal detachment and coronary artery disease. Alternative splicing results in multi ple transcript variants.

基因ID	9079
基因名	LDB2
Swiss	O43679
别名	LDB2;CLIM1;LDB-2;LDB1

产品验证



Western blot analysis of LDB2 expressed in U-87MG,Raji,Mouse kidney using LDB2 Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Block ing buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn