

MAN2B1 Rabbit pAb

货号**: B15705**

产品信息

nal
:500 - 1:2000
t -20°C. Avoid freeze / thaw cycles. PBS with 0.02% sodium azide,50% glycerol,pH7.3.
ugated
liver
me
purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 640-850 of human MAN 2B1 (NP_000519.2).
序列	TFFWYNASIGDNESDQASGAYIFRPNQQKPLPVSRWAQIHLVKTPLVQEVHQNFSAWCSQVVRLYPGQRHLELEWSVGP IPVGDTWGKEVISRFDTPLETKGRFYTDSNGREILERRRDYRPTWKLNQTEPVAGNYYPVNTRIYITDGNMQLTVLTDRSQ GGSSLRDGSLELMVHRRLLKDDGRGVSEPLMENGSGAWVRGRHLVLLDTAQ

靶点信息

研究背景	This gene encodes an enzyme that hydrolyzes terminal, non-reducing alpha-D-mannose residues in alpha -D-mannosides. Its activity is necessary for the catabolism of N-linked carbohydrates released during glyc oprotein turnover and it is member of family 38 of glycosyl hydrolases. The full length protein is processe d in two steps. First, a 49 aa leader sequence is cleaved off and the remainder of the protein is processed into 3 peptides of 70 kDa, 42 kDa (D) and 13/15 kDa (E). Next, the 70 kDa peptide is further processed in to three peptides (A, B and C). The A, B and C peptides are disulfide-linked. Defects in this gene have been associated with lysosomal alpha-mannosidosis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
基因ID	4125
基因名	MAN2B1
Swiss	000754
别名	MAN2B1;LAMAN;MANB

产品验证



Western blot analysis of MAN2B1 expressed in Mouse liver using MAN2B1 Rabbit pAb at 1:1000. Secondar y antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn