

ATP5L Rabbit pAb

货号: B15831

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 11kDa |
| 实测分子量 | 11kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | 293T,A-549,HepG2,HeLa,Mouse brain,Mouse liver,Rat brain,Rat kidney |
| 细胞定位 | Mitochondrion,Mitochondrion inner membrane |
| 纯化 | Affinity purification |

抗原信息

| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-103 of human ATP5L (NP_006467.4). |
| 序列 | MAQFVRNLVEKTPALVNAAVTYSKPRLATFWYYYAKVELVPPTPAEIPRAIQSLKKIVNSAQTGSKQLTVKEAVLNGLVATEV LMWFYVGEIIGKRGIIGYDV |


靶点信息

| | |
|------|--|
| 研究背景 | Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, which comprises the proton channel. The F1 complex consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled in a ratio of 3 alpha, 3 beta, and a single representative of the other 3. The Fo seems to have nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene encodes the g subunit of the Fo complex. Alternative splicing results in multiple transcript variants. |
|------|--|

| | |
|-------|--------------|
| 基因ID | 10632 |
| 基因名 | ATP5L |
| Swiss | O75964 |
| 别名 | ATP5L;ATP5JG |

产品验证

Western blot analysis of ATP5L expressed in Mouse brain,Rat brain,293T using ATP5L Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn