

OPRK1 Rabbit pAb

货号: **B16042**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	33kDa/42kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,HeLa,U-937,Mouse brain,Rat brain
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-60 of human OPRK1 (NP_000903.2).
序列	MDSPIQIFRGEPGPTCAPSACLPPNSSAWFPGWAEPDSNGSAGSEDAQLEPAHISPAIPV

靶点信息

研究背景	This gene encodes an opioid receptor, which is a member of the 7 transmembrane-spanning G protein-co upled receptor family. It functions as a receptor for endogenous ligands, as well as a receptor for various synthetic opioids. Ligand binding results in inhibition of adenylate cyclase activity and neurotransmitter r elease. This opioid receptor plays a role in the perception of pain and mediating the hypolocomotor, analg esic and aversive actions of synthetic opioids. Variations in this gene have also been associated with alco hol dependence and opiate addiction. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. A recent study provided evidence for translational readthrough in this gene and expression of an additional C-terminally extended isoform via the use of an alternative in-frame translation termination codon.
基因 ID	4986
基因名	OPRK1
Swiss	P41145
别名	OPRK1;K-OR-1;KOR;KOR-1;OPRK

产品验证



Western blot analysis of OPRK1 expressed in Mouse brain,Rat brain,U-251MG using OPRK1 Rabbit pAb a t 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per la ne. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn