

CHRFAM7A Rabbit pAb

货号: B16189

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	46kDa
实测分子量	46kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,Jurkat,SH-SY5Y,U-87MG,NIH/3T3,Mouse brain,Mouse spinal cord,Rat brain,Rat spinal cord
细胞定位	Membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-150 of human CHRFA M7A (NP_647536.1).
序列	MQKYCIYQHFQFQLLIQHLWIAANCDIADERFDATFHTNVLVNSSGHCQYLPPGIFKSSCYIDVRWFPPFDVQHCKLKFGS WSYGGWSLDLQMQEADISGYIPNGEWDLVGIPGKRSEFYECCKEYPDPVTFTVTMRRRTLYYGLNLLIP

靶点信息

研究背景	The nicotinic acetylcholine receptors (nAChRs) are members of a superfamily of ligand-gated ion channels that mediate fast signal transmission at synapses. The family member CHRNA7, which is located on chromosome 15 in a region associated with several neuropsychiatric disorders, is partially duplicated and forms a hybrid with a novel gene from the family with sequence similarity 7 (FAM7A). Alternative splicing has been observed, and two variants exist, for this hybrid gene. The N-terminally truncated products predicted by the largest open reading frames for each variant would lack the majority of the neurotransmitter-gated ion-channel ligand binding domain but retain the transmembrane region that forms the ion channel. Although current evidence supports transcription of this hybrid gene, translation of the nicotinic acetylcholine receptor-like protein-encoding open reading frames has not been confirmed.
基因ID	89832
基因名	CHRFAM7A
Swiss	Q494W8
别名	CHRFAM7A;CHRNA7;CHRNA7-DR1;D-10

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn