

GSTM3 Rabbit pAb

货号: **B16243**

产品信息

反应	Human, Mouse, Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	26kDa
实测分子量	25kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,Mouse liver,Mouse testis
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 101-225 of human GST M3 (NP_000840.2).
序列	VDIIENQVMDFRTQLIRLCYSSDHEKLKPQYLEELPGQLKQFSMFLGKFSWFAGEKLTFVDFLTYDILDQNRIFDPKCLDEF PNLKAFMCRFEALEKIAAYLQSDQFCKMPINNKMAQWGNKPVC

靶点信息

研究背景	Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergen e families. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transfer ases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. This gene encodes a gluta thione S-transferase that belongs to the mu class. The mu class of enzymes functions in the detoxificatio n of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and produc ts of oxidative stress, by conjugation with glutathione. The genes encoding the mu class of enzymes are organized in a gene cluster on chromosome 1p13.3 and are known to be highly polymorphic. These genet ic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxici ty and efficacy of certain drugs. Mutations of this class mu gene have been linked with a slight increase in a number of cancers, likely due to exposure with environmental toxins. Alternative splicing results in multiple transcript variants.
基因ID	2947
基因名	GSTM3
Swiss	P21266
别名	GSTM3;GST5;GSTB;GSTM3-3;GTM3

产品验证



Western blot analysis of GSTM3 expressed in Mouse liver, HeLa, HepG2 using GSTM3 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn