

# PCDHA13 Rabbit pAb

货号: **B16324** 

### 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	87kDa/102kDa
实测分子量	90kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Mouse kidney,Rat brain,Rat kidney
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


### 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 300-360 of human PCD HA13 (NP_061727.1).
序列	GEIRTKGKLDFEEKKLYEISVEAVDKGNIPMAGHCTLLVEVLDVNDNAPEVTITSLSLPIR

靶点信息

研究背景	This gene is a member of the protocadherin alpha gene cluster, one of three related gene clusters tande mly linked on chromosome five that demonstrate an unusual genomic organization similar to that of B-cel I and T-cell receptor gene clusters. The alpha gene cluster is composed of 15 cadherin superfamily genes related to the mouse CNR genes and consists of 13 highly similar and 2 more distantly related coding seq uences. The tandem array of 15 N-terminal exons, or variable exons, are followed by downstream C-termi nal exons, or constant exons, which are shared by all genes in the cluster. The large, uninterrupted N-ter minal exons each encode six cadherin ectodomains while the C-terminal exons encode the cytoplasmic d omain. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins that mo st likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been observed and additional variants have been suggested but their full-length nature has yet to be determined.
基因ID	56136
基因名	PCDHA13
Swiss	Q9Y5I0
别名	PCDHA13;CNR5;CNRN5;CNRS5;CRNR5;PCDH-ALPHA13

#### 产品验证



Western blot analysis of PCDHA13 expressed in Mouse brain,Rat brain,Mouse kidney using PCDHA13 R abbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure ti me: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn