

# CYP2C18 Rabbit pAb

货号: B16344

### 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	48kDa/55kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver,Rat liver
细胞定位	Endoplasmic reticulum membrane,Microsome membrane,Peripheral membrane protein
纯化	Affinity purification


### 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 231-490 of human CYP2 C18 (NP_000763.1).
序列	NKIAENFAYIKSYVLERIKEHQESLDMNSARDFIDCFLIKMEQEKHNQQSEFTVESLIATVTDMFGAGTETTSTTLRYGLLLLL KYPEVTAKVQEEIECVVGRNRSPCMQDRSHMPYTDAVVHEIQRYIDLLPTNLPHAVTCDVKFKNYLIPKGTTIITSLTSVLHN DKEFPNPEMFDPGHFLDKSGNFKKSDYFMPFSAGKRMCMGEGLARMELFLFLTTILQNFNLKSQVDPKDIDITPIANAFGR VPPLYQLCFIPV

靶点信息

研究背景	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 pro teins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of c holesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum but its specific su bstrate has not yet been determined. The gene is located within a cluster of cytochrome P450 genes on c hromosome 10q24. An additional gene, CYP2C17, was once thought to exist; however, CYP2C17 is now c onsidered an artefact based on a chimera of CYP2C18 and CYP2C19. Alternatively spliced transcript varia nts encoding different isoforms have been found for this gene.
基因ID	1562
基因名	CYP2C18
Swiss	P33260
别名	CYP2C18;CPCI;CYP2C;CYP2C17;P450-6B/29C;P450IIC17

### 产品验证



Western blot analysis of CYP2C18 expressed in Mouse liver, Rat liver using CYP2C18 Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit lgG(H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn