

ACAA1 Rabbit pAb

货号: B16349

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	34kDa/44kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HL-60,MCF7,A-549,Jurkat
细胞定位	Peroxisome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 27-300 of human ACAA1 (NP_001598.1).
序列	LSGAPQASAADV VVHGRRTAICRAGRGGFKDTTPDELLSAVMTAVLKDVNLRPEQLGDCVGNVLQPGAGAIMARIAQF LSDIPETVPLSTVNRQCSSGLQAVASIAGGIRNGSYDIGMACGVESMSLADRGNPGNITSRLMEKEKARDCLIPMGITSEN VAERFGISREKQDTFALASQQKAARAQSKGCFQAEIVPVT TTVHDDKGTKRSITVTQDEGIRPSTTMEGLAKLKPAFKKDG STTAGNSSQVSDGAAAILARRSKAEELGLPI

靶点信息

研究背景	This gene encodes an enzyme operative in the beta-oxidation system of the peroxisomes. Deficiency of th is enzyme leads to pseudo-Zellweger syndrome. Alternative splicing results in multiple transcript variants .
基因ID	30

基因名	ACAA1
Swiss	P09110
别名	ACAA1;ACAA;PTHIO;THIO

产品验证



Western blot analysis of ACAA1 expressed in HL-60,MCF7,A-549 using ACAA1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn