

MTF2 Rabbit pAb

货号**: B16485**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	29kDa/55kDa/60kDa/67kDa
实测分子量	67kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HL-60,HepG2,Mouse brain,Rat brain
细胞定位	Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-190 of human MTF2 (NP_001157864.1).
序列	MRDSTGAGNSLVHKRSPLRRNQKTPTSLTKLSLQDGHKAKKPACKFEEGQDVLARWSDGLFYLGTIKKINILKQSCFIIFE DSSKSWVLWKDIQTGATGSGEMVCTICQEEYSEAPNEMVICDKCGQGYHQLCHTPHIDSSVIDSDEKWLCRQCVFATTT KRGGALKKGPNAKALQVMKQTLPYSVADLE

靶点信息

研究背景	Polycomb group (PcG protein that specifically binds histone H3 trimethylated at 'Lys-36' (H3K36me3 and recruits the PRC2 complex, thus enhancing PRC2 H3K27me3 methylation activity. Regulates the transcrip tional networks during embryonic stem cell self-renewal and differentiation (By similarity. Promotes recru itment of the PRC2 complex to the inactive X chromosome in differentiating XX ES cells and PRC2 recruit ment to target genes in undifferentiated ES cells (By similarity. Required to repress Hox genes by enhanc ing H3K27me3 methylation of the PRC2 complex (By similarity. In some conditions may act as an inhibito r of PRC2 activity: able to activate the CDKN2A gene and promote cellular senescence by suppressing the catalytic activity of the PRC2 complex locally (By similarity. Binds to the metal-regulating-element (MRE o f MT1A gene promoter (By similarity.
基因ID	22823
基因名	MTF2
Swiss	Q9Y483
别名	MTF2;M96;PCL2;TDRD19A;dJ976O13.2

产品验证



Western blot analysis of MTF2 expressed in Mouse brain,Rat brain,HL-60 using MTF2 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per Iane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn