

EIF2B2 Rabbit pAb

货号: B16507

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:50 - 1:200
理论分子量	38kDa
实测分子量	39kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse thymus,Rat kidney
细胞定位	cytoplasm,cytosol,eukaryotic translation initiation factor 2B complex
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-120 of human EIF2B2 (NP_055054.1).
序列	MPGSAAKGSELSESFVETLKRGGGPRSSSEEMARETLGLLRQIITDHRWSNAGELMELIRREGRRMTAAQPSETTVGNM VRRVLKIIREEYGR LHGRSDESDQQESLHKLLTSGGLNE

靶点信息

研究背景	This gene encodes the beta subunit of eukaryotic initiation factor-2B (EIF2B). EIF2B is involved in protein synthesis and exchanges GDP and GTP for its activation and deactivation.
基因ID	8892

基因名	EIF2B2
Swiss	P49770
别名	EIF2B2;EIF-2Bbeta;EIF2B

产品验证



Western blot analysis of EIF2B2 expressed in Mouse thymus,Rat kidney using EIF2B2 Rabbit pAb at 1:100 0. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn