

PDE5A Rabbit pAb

货号: B16583

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	94kDa/99kDa
实测分子量	100kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-1080,SKOV3,A-549,22Rv1,Mouse brain,Mouse skeletal muscle,Rat skeletal muscle
细胞定位	cytosol
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-150 of human PDE5A (NP_001074.2).
序列	MERAGPSFGQQRQQQPQQKQQQRDQDSVEAWLDDHWDFTFSYFVRKATREMVNAWFAERVHTIPVCKEGIRGHT ESCSCPLQQSPRADNSAPGTPTRKISASEFDRPLRPVVKDSEGTVSFLSDSEKKEQMPLTPPRFDHDEGDQCS

靶点信息

研究背景	This gene encodes a cGMP-binding, cGMP-specific phosphodiesterase, a member of the cyclic nucleotide phosphodiesterase family. This phosphodiesterase specifically hydrolyzes cGMP to 5'-GMP. It is involved in the regulation of intracellular concentrations of cyclic nucleotides and is important for smooth muscle relaxation in the cardiovascular system. Alternative splicing of this gene results in three transcript variants encoding distinct isoforms.
基因ID	8654

基因名	PDE5A
Swiss	O76074
别名	PDE5A;CGB-PDE;CN5A;PDE5

产品验证



Western blot analysis of PDE5A expressed in Mouse brain,Rat skeletal muscle,HT-1080 using PDE5A Rab bit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30u g per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

实验步骤

访问官网浏览详情: www.ablybio.cn