

RAD54L2 Rabbit pAb

货号: B16838

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	162kDa
实测分子量	162kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,THP-1
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1178-1467 of human RA D54L2 (NP_055921.2).
序列	ELQQYADVAAARESQRQSSPSTNAALPGPPAQLMDSSAVPGTALGTEPRLGGHCLNSSLLVTGQPCGDRHPVLDLRGHKR KLATPPAAQESSRRRSRKGHLPAPVQPYEHGYPVSGGFAMPPVSLNHNLTTPFTSQAGENSLFMGSTPSYYQLSNLLADA RLVFPVTTDPLVPAGPVSSSSTATSVTASNPSFMLNPSPVGPILPSYSLPFSQPLLSEPRMFAPFPSPVLPSNLSRGMSIYPGY MSPHAGYPAGLLRSQVPPFDSHEVAEVGFSSNDDDEDKDDDVIEVTGK

靶点信息

研究背景	DNA helicase that modulates androgen receptor (AR-dependent transactivation in a promoter-dependent manner. Not able to remodel mononucleosomes in vitro (By similarity.
基因ID	23132

基因名	RAD54L2
Swiss	Q9Y4B4
别名	RAD54L2;ARIP4;HSPC325;SRISNF2L

产品验证



Western blot analysis of RAD54L2 expressed in MCF7,THP-1 using RAD54L2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn