

CD22 Rabbit pAb

货号: B16919

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	75kDa/84kDa/85kDa/95kDa
实测分子量	140KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse thymus,Rat lung
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 708-847 of human CD22 (NP_001762.2).
序列	LQRRWKRTQSQQGLQENSSGQSFFVRNKKVRRAPLSEGPLSLGCYNPMMEDGISYTTLRFPENIPRTGDAESSEMQR PPPCDDTVTYSALHKRQVGDYENVIPDFPEDEGIHYSELIQFGVGERPQAQENVDYVILKH

靶点信息

研究背景	Siglecs (sialic acid binding Ig-like lectins) are I-type (Ig-type) lectins belonging to the Ig superfamily. They are characterized by an N-terminal Ig-like V-type domain which mediates sialic acid binding, followed by varying numbers of Ig-like C2-type domains. Human Siglec-2, also known as B-cell antigen CD22 or Blymphocyte cell adhesion molecule (BL-CAM), is a B-cell restricted glycoprotein that is expressed in the cytoplasm of progenitor B and pre-B cells and on the surface of mature B cells. Two distinct human Siglec2/CD22 cDNAs that arise from differential RNA processing of the same gene have been isolated. Siglec2/CD22 is an adhesion molecule that preferentially binds alpha 2,6- linked sialic acid on the same (cis) or adjacent (trans) cells. Interaction of CD22 with trans ligands on opposing cells was found to be favored over the binding of ligands in cis.
基因ID	933
基因名	CD22
Swiss	P20273
别名	CD22;SIGLEC-2;SIGLEC2

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn