

PCGF6 Rabbit pAb

货号: **B16931**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	30kDa/39kDa
实测分子量	39kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 161-350 of human PCGF 6 (NP_001011663.1).
序列	RHFYYSNRCPKCNIVVHQTQPLYNIRLDRQLQDIVYKLVINLEEREKKQMHDFYKERGLEVPKPAVPQPVPSSKGRSKKVLE SVFRIPPELDMSLLLEFIGANEGTGHFKPLEKKFVRVSGEATIGHVEKFLRRKMGLDPACQVDIICGDHLLEQYQTLREIRRAI GDAAMQDGLLVLHYGLVVSPLKIT

靶点信息

研究背景	The protein encoded by this gene contains a RING finger motif, which is most closely related to those of p olycomb group (PcG) proteins RNF110/MEL-18 and BMI1. PcG proteins are known to form protein complex es and function as transcription repressors. This protein has been shown to interact with some PcG proteins and act as a transcription repressor. The activity of this protein is found to be regulated by cell cycle dependent phosphorylation. Alternatively spliced transcript variants encoding different isoforms have been identified.
------	---

基因 ID	84108
基因名	PCGF6
Swiss	Q9BYE7
别名	PCGF6;MBLR;RNF134

产品验证



Western blot analysis of PCGF6 expressed in MCF7 using PCGF6 Rabbit pAb at 1:1000. Secondary antibod y: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn