

# PCSK5 Rabbit pAb

货号: B17037

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	101kDa/206kDa
实测分子量	102kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,SW480,Mouse lung,Rat lung,Rat brain
细胞定位	Endomembrane system,Secreted,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 601-913 of human PCSK 5 (NP_006191.2).
序列	QPYSPTNEFPKVERFRYSRVEDPTDDYGTEDYAGPCDPECSEVGCDGPGPDHCNDCLHYYYKLKNNTRICVSSCPPGHYH ADKKRCRKCAPNCESCFGSHGDQCMSCKYGYFLNEETNSCVTHCPDGSYQDTKKNLCRKSENCKTCTEFHNCTECRD GLSLQGSRCSVSCEDGRYFNGQDCQPCHRFCATCAGAGADGCINCTEGYFMEDGRCVQSCSISYYFDHSSSENGYKSCK KCDISCLTCNGPGFKNCTSCPSGYLLDLGMCQMGAICKDATEESWAEGGFCMLVKNNLQCRKVLQQLCCKTCTFQG

靶点信息

研究背景	This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an initial autocatalytic processing event in the ER to generate a heterodimer which exits the ER. It then sorts to the trans-Golgi network where a second autocatalytic event takes place and the catalytic activity is acquired. This encoded protein is widely expressed and one of the seven basic amino acid-specific members which cleave their substrates at single or paired basic residues. It mediates posttranslational endoproteolytic processing for several integrin alpha subunits and is thought to process prorenin, pro-membrane type-1 matrix metalloproteinase and HIV-1 glycoprotein gp160. Alternative splicing results in multiple transcript variants, some of which encode distinct isoforms, including a protease packaged into dense core granules (PC5A) and a type 1 membrane bound protease (PC5B).
基因ID	5125
基因名	PCSK5
Swiss	Q92824
别名	PCSK5;PC5;PC6;PC6A;SPC6

产品验证



Western blot analysis of PCSK5 expressed in Mouse lung,Rat lung,Jurkat using PCSK5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)