

TriMethyl-Histone H3-K14 Rabbit pAb

货号: **B17100**

产品信息

| 反应 | Human, Mouse, Rat |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IHC IP CHIP |
| 推荐浓度 | WB: 1:100 - 1:500 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:50 - 1:200 ChIP: 1:20 - 1:100 |
| 理论分子量 | 15kDa |
| 实测分子量 | 17kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,NIH/3T3 |
| 细胞定位 | Chromosome, Nucleus |
| 纯化 | Affinity purification |


抗原信息

| 抗原信息 | A synthetic trimethylated peptide around K14 of human Histone H3 (NP_003520.1). |
|------|---|
| 序列 | GGKAP |

靶点信息

| 研究背景 | Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octa mer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent his tone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3. |
|--------------|---|
| 基因 ID | 8290 |
| 基因名 | HIST3H3 |
| Swiss | Q16695 |
| 别名 | H3.4;H3/g;H3FT;H3t;HIST3H3;Histone H3;HIST1H3A |

产品验证



Western blot analysis of TriMethyl-Histone H3-K14 expressed in HeLa,NIH/3T3 using TriMethyl-Histone H3-K14 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. L ysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn