

CLCNKA Rabbit pAb

货号**: B17279**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	70kDa/75kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,HT-29,A-549
细胞定位	Membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 470-644 of human CLCN KA (NP_001244068.1).	
序列	QSCQPSFYDGTIIVKKLPYLPRILGRNIGSHHVRVEHFMNHSITTLAKDTPLEEVVKVVTSTDVTEYPLVESTESQILVGIVQR AQLVQALQAEPPSRAPGHQQCLQDILARGCPTEPVTLTLFSETTLHQAQNLFKLLNLQSLFVTSRGRAVGCVSWVEMKKAI SNLTNPPAPK	

靶点信息

研究背景	This gene is a member of the CLC family of voltage-gated chloride channels. The encoded protein is predicted to have 12 transmembrane domains, and requires a beta subunit called barttin to form a functional channel. It is thought to function in salt reabsorption in the kidney and potassium recycling in the inner e ar. The gene is highly similar to CLCNKB, which is located 10 kb downstream from this gene. Multiple transcript variants encoding different isoforms have been found for this gene.

基因ID	1187
基因名	CLCNKA
Swiss	P51800
别名	CLCNKA;CLCK1;CIC-K1;hCIC-Ka

产品验证



Western blot analysis of CLCNKA expressed in LO2,HT-29,A-549 using CLCNKA Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bloc king buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn