

CETN1 Rabbit pAb

货号: B17284

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 19kDa |
| 实测分子量 | 20kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | 293T,Jurkat,MCF7,THP-1,HeLa,Mouse testis,Rat testis |
| 细胞定位 | Cytoplasm,centrosome,cytoskeleton,microtubule organizing center |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-172 of human CETN1 (NP_004057.1). |
| 序列 | MASGFKKPSAAGTGQKRKVAPKPELTEDQKQEVRSEAFDLFDVGSGTIDAKELKVAMRALGFEPRKEMKKMISEVDRE GTGKISFNDFLAVMTQKMSEKDTKEEILKAFLRFDDDETGKISFKNLKRVANELGENLTDEELQEMIDEADRDRGDGEVNE EEFLRIMKKTSY |

靶点信息

| | |
|------|--|
| 研究背景 | The protein encoded by this gene plays important roles in the determination of centrosome position and segregation, and in the process of microtubule severing. This protein is localized to the centrosome of interphase cells, and redistributes to the region of the spindle poles during mitosis, reflecting the dynamic behavior of the centrosome during the cell cycle. |
| 基因ID | 1068 |

| | |
|-------|---------------------------|
| 基因名 | CETN1 |
| Swiss | Q12798 |
| 别名 | CETN1;CEN1;CETN;centrin-1 |

产品验证



Western blot analysis of CETN1 expressed in Mouse testis, Rat testis, 293T using CETN1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn