

SUMO3 Rabbit pAb

货号: **B17370**

产品信息

反应 Human,Mouse,Rat 和主 Rabbit 文陰性 Polyclonal 例測反应 应用 WB IHC 推荐秩度 WB: 1:1000 - 1:2000 IHC: 1:50 - 1:100 理论分子嚴 11kDa/15kDa 支剥分子量 19kDa 形式 Liquid 保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. 何戰物 Unconjugated 阳性对照 MCF7 细胞定位 Cytoplasm,Nucleus,PML body		
克隆性 Polyclonal 預測反应 WB IHC 推荐浓度 WB: 1:1000 - 1:2000 IHC: 1:50 - 1:100 理论分子量 11kDa/15kDa 实测分子量 19kDa 形式 Liquid 保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. 俱联物 Unconjugated 阿性对照 MCF7	反应	Human,Mouse,Rat
	宿主	Rabbit
应用 WB IHC 推荐來度 WB: 1:1000 - 1:2000 IHC: 1:50 - 1:100 理论分子量 11kDa/15kDa 实測分子量 19kDa 形式 Liquid 保存条件 Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. 佛联物 Unconjugated 阳性对照 MCF7	克隆性	Polyclonal
# # # # # # # # # # # # # # # # # # #	预测反应	
IHC: 1:50 - 1:100 理论分子量	应用	WB IHC
实测分子量 19kDa 形式 Liquid 保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. 偶联物 Unconjugated 阳性对照 MCF7	推荐浓度	
形式 Liquid 保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. (周联物 Unconjugated 阳性对照 MCF7	理论分子量	11kDa/15kDa
保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. (周联物 Unconjugated MCF7	实测分子量	19kDa
Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. (周联物 Unconjugated MCF7	形式	Liquid
阳性对照 MCF7	保存条件	
	偶联物	Unconjugated
细胞定位 Cytoplasm,Nucleus,PML body	阳性对照	MCF7
	细胞定位	Cytoplasm,Nucleus,PML body
纯化 Affinity purification	纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 40-103 of human SUMO3 (P55854).
序列	SKLMKAYCERQGLSMRQIRFRFDGQPINETDTPAQLEMEDEDTIDVFQQQTGGVPESSLAGHSF

靶点信息

研究背景	This gene encodes a member of the small ubiquitin-related modifier (SUMO) family of eukaryotic proteins. The encoded protein is covalently conjugated to other proteins via a post-translation modification known as sumoylation. Sumoylation may play a role in a wide variety of cellular processes, including nuclear transport, DNA replication and repair, mitosis, transcriptional regulation, and signal transduction. Alternative ly spliced transcript variants encoding distinct proteins have been described.
基因 ID	6612

基因名	SUMO3
Swiss	P55854
别名	SUMO3;SMT3A;SMT3H1;SUMO-3;Smt3B

产品验证



Western blot analysis of SUMO3 expressed in MCF7 using SUMO3 Rabbit pAb at 1:1000. Secondary antibo dy: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-f at dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn