

MC1 Receptor Rabbit pAb

货号: B17394

产品信息

反应	Human, Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:1000 - 1:2000
理论分子量	34kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	M21,MCF7,SW620,22Rv1,Raji,Mouse testis
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant protein of human MC1 Receptor
序列	

靶点信息

研究背景	This intronless gene encodes the receptor protein for melanocyte-stimulating hormone (MSH). The encod ed protein, a seven pass transmembrane G protein coupled receptor, controls melanogenesis. Two types of melanin exist: red pheomelanin and black eumelanin. Gene mutations that lead to a loss in function ar e associated with increased pheomelanin production, which leads to lighter skin and hair color. Eumelani n is photoprotective but pheomelanin may contribute to UV-induced skin damage by generating free radi cals upon UV radiation. Binding of MSH to its receptor activates the receptor and stimulates eumelanin sy nthesis. This receptor is a major determining factor in sun sensitivity and is a genetic risk factor for melan oma and non-melanoma skin cancer. Over 30 variant alleles have been identified which correlate with ski n and hair color, providing evidence that this gene is an important component in determining normal hu man pigment variation.
基因 ID	4157
基因名	MC1R
Swiss	Q01726
别名	MC1R;CMM5;MSH-R;SHEP2

产品验证



Western blot analysis of MC1 Receptor expressed in Mouse testis,M21,MCF7 using MC1 Receptor Rab bit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure ti me: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn