

# **GNAS Rabbit pAb**

货号: **B17463** 

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	28kDa/44kDa/45kDa/77kDa/109kDa/111kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NCI-H460,SW620,Mouse skeletal muscle,Mouse brain,Mouse heart
细胞定位	Cell membrane By similarity; Lipid-anchor By similarity
纯化	Affinity purification


# 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human GNAS (P63092).
序列	MGCLGNSKTEDQRNEEKAQREANKKIEKQLQKDKQVYRATHRLLLLGAGESGKSTIVKQMRILHVNGFNGEGGEEDPQA ARSNSDGEKATKVQDIKNNLK

靶点信息

研究背景	This locus has a highly complex imprinted expression pattern. It gives rise to maternally, paternally, and biallelically expressed transcripts that are derived from four alternative promoters and 5' exons. Some tra nscripts contain a differentially methylated region (DMR) at their 5' exons, and this DMR is commonly fou nd in imprinted genes and correlates with transcript expression. An antisense transcript is produced from an overlapping locus on the opposite strand. One of the transcripts produced from this locus, and the anti sense transcript, are paternally expressed noncoding RNAs, and may regulate imprinting in this region. In addition, one of the transcripts contains a second overlapping ORF, which encodes a structurally unrelate d protein - Alex. Alternative splicing of downstream exons is also observed, which results in different form s of the stimulatory G-protein alpha subunit, a key element of the classical signal transduction pathway li nking receptor-ligand interactions with the activation of adenylyl cyclase and a variety of cellular reponse s. Multiple transcript variants encoding different isoforms have been found for this gene. Mutations in this gene result in pseudohypoparathyroidism type 1a, pseudohypoparathyroidism type 1b, Albright hereditar y osteodystrophy, pseudopseudohypoparathyroidism, McCune-Albright syndrome, progressive osseus het eroplasia, polyostotic fibrous dysplasia of bone, and some pituitary tumors.
基因 <b>ID</b>	2778
基因名	GNAS
Swiss	O95467,P63092,P84996,Q5JWF2
别名	GNAS;AHO;C20orf45;GNAS1;GPSA;GSA;GSP;NESP;POH;SCG6;SgVI

## 产品验证



Western blot analysis of GNAS expressed in Mouse skeletal muscle, NCI-H460, SW620 using GNAS Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug p er lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

#### 实验步骤

访问官网浏览详情: www.ablybio.cn