

NEDD9 Rabbit pAb

货号: B17506

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	19kDa/92kDa
实测分子量	105KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549
细胞定位	Cell junction,Cell projection,Cytoplasm,Golgi apparatus,Nucleus,cell cortex,cytoskeleton,focal adhesion,la mellipodium,spindle
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 390-560 of human NED D9 (NP_006394.1).
序列	SSLSASPAQDKRLFLDPDTAIERLQRLQQALEMGVSSLMALVTTDWRCYGYMERHINEIRTAVDKVELFLKEYLHFVKGAV ANAACLP ELILHNKMKRELQRVEDSHQILSQTSHDLNECSWSLNILAINKPQNKCDLDRFVMVAKTVPDDAKQLTTTINT NAEALFRPG

靶点信息

研究背景	The protein encoded by this gene is a member of the CRK-associated substrates family. Members of this family are adhesion docking molecules that mediate protein-protein interactions for signal transduction pathways. This protein is a focal adhesion protein that acts as a scaffold to regulate signaling complexes important in cell attachment, migration and invasion as well as apoptosis and the cell cycle. This protein has also been reported to have a role in cancer metastasis. Alternative splicing results in multiple transcript variants.
基因ID	4739
基因名	NEDD9
Swiss	Q14511
别名	NEDD9;CAS-L;CAS2;CASL;CASS2;HEF1

产品验证



Western blot analysis of NEDD9 expressed in A-549 using NEDD9 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn