

MT-ATP6 Rabbit pAb

货号: B17512

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	25kDa
实测分子量	25kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	mitochondrial inner membrane,mitochondrial proton-transporting ATP synthase complex
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 50-150 of human MT-ATP6 (YP_003024031.1).
序列	IKLTSKQMMTMHNTKGRTWSMLVSLIIFIATTNLLGLPHSFTPTTQLSMNLAMAIPWAGTVIMGFRSKIKNALAHFLPQG TPTPLIPMLVIIETISLL

靶点信息

研究背景	Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembranous catalytic core and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Key component of the proton channel; it may play a direct role in the translocation of protons across the membrane.
基因ID	4508
基因名	MT-ATP6
Swiss	P00846
别名	ATPase6;MTATP6;MT-ATP6

产品验证

Western blot analysis of MT-ATP6 expressed in mouse lung, rat liver using MT-ATP6 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn