

CBX1 Rabbit pAb

货号: **B17549**

产品信息

| 反应 | Human |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:1000 |
| 理论分子量 | 21kDa |
| 实测分子量 | 21kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | OVCAR3 |
| 细胞定位 | Nucleus |
| 纯化 | Affinity purification |


抗原信息

| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 1-100 of human HP1 beta/CBX1 (NP_ 006798.1). | |
|------|---|--|
| 序列 | MGKKQNKKKVEEVLEEEEEEYVVEKVLDRRVVKGKVEYLLKWKGFSDEDNTWEPEENLDCPDLIAEFLQSQKTAHETDKS EGGKRKADSDSEDKGEESKP | |

靶点信息

| 研究背景 | This gene encodes a highly conserved nonhistone protein, which is a member of the heterochromatin prot ein family . The protein is enriched in the heterochromatin and associated with centromeres. The protein has a single N-terminal chromodomain which can bind to histone proteins via methylated lysine residues, and a C-terminal chromo shadow-domain (CSD) which is responsible for the homodimerization and interaction with a number of chromatin-associated nonhistone proteins. The protein may play an important role in the epigenetic control of chromatin structure and gene expression. Several related pseudogenes are located on chromosomes 1, 3, and X. Multiple alternatively spliced variants, encoding the same protein, have been identified. |
|-------|---|
| 基因ID | 10951 |
| 基因名 | CBX1 |
| Swiss | P83916 |
| 别名 | CBX1;CBX;HP1-BETA;HP1Hs-beta;HP1Hsbeta;M31;MOD1;p25beta |

产品验证



Western blot analysis of CBX1 expressed in OVCAR3 using CBX1 Rabbit pAb at 1:1000. Secondary antibod y: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn