

DUSP2 Rabbit pAb

货号: **B17590**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	34kDa
实测分子量	34kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,NIH/3T3,HeLa,MCF7
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 150-250 of human DUSP2 (NP_0044 09.1).	
序列	ALPPTGDKTSRSDSRAPVYDQGGPVEILPYLFLGSCSHSSDLQGLQACGITAVLNVSASCPNHFEGLFRYKSIPVEDNQMV EISAWFQEAIGFIDWVKNSG	

靶点信息

研究背景	The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. Thes e phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine a nd phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate s pecificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK1 and ERK2, is predominantly expressed in hematopoietic tissues, and is localized in the nucleus.
基因ID	1844
基因名	DUSP2
Swiss	Q05923
别名	DUSP2;PAC-1;PAC1

产品验证



Western blot analysis of DUSP2 expressed in 293T,NIH/3T3,HeLa using DUSP2 Rabbit pAb at 1:1000. S econdary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocki ng buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn