

MTIF2 Rabbit pAb

货号: B17613

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100
理论分子量	81kDa
实测分子量	81kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,A-431,U-251MG,Mouse heart,Mouse kidney,Mouse liver,Mouse spleen,Rat testis
细胞定位	Mitochondrion
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 428-727 of human MTIF 2 (NP_002444.2).
序列	GEEILEVESEPRAREVVDWRKYEQEKEKGQEDLKIIIEEKKEHKEAHQKAREKYGHLLWKKRSILRFLERKEQIPLKPKEKRE RDSNVLSVIIKGDVDGGSVEAILNIIDTYDASHECELELVHFGVGDVSANDVNLAETFDGVIYGFNVNAGNVIQQSAAKKG V KIKLHKIYRLVEDLQEELSSRLPCAVEEHPVGEASILATFSVTEGKKKVPVAGCRVQKGQLEKQKKFKLTRNGHVIWKGSLT SLKHHKDDISIVKTGMDCGLSLDEDNMEFQVGDRIVCYEEKQIQAKTSWDPGF

靶点信息

研究背景	During the initiation of protein biosynthesis, initiation factor-2 (IF-2) promotes the binding of the initiator tRNA to the small subunit of the ribosome in a GTP-dependent manner. Prokaryotic IF-2 is a single polypeptide, while eukaryotic cytoplasmic IF-2 (eIF-2) is a trimeric protein. Bovine liver mitochondria contain IF-2(mt), an 85-kD monomeric protein that is equivalent to prokaryotic IF-2. The predicted 727-amino acid human protein contains a 29-amino acid presequence. Human IF-2(mt) shares 32 to 38% amino acid sequence identity with yeast IF-2(mt) and several prokaryotic IF-2s, with the greatest degree of conservation in the G domains of the proteins.
基因ID	4528
基因名	MTIF2
Swiss	P46199
别名	MTIF2

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn