

TNFSF12 Rabbit pAb

货号**: B17651**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100
理论分子量	27kDa/36kDa
实测分子量	27kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat kidney
细胞定位	Cell membrane,Secreted,Single-pass membrane protein,Single-pass type II membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 42-123 of human TNFSF 12 (O43508).	
序列	VSLGSRASLSAQEPAQEELVAEEDQDPSELNPQTEESQDPAPFLNRLVRPRRSAPKGRKTRARRAIAAHYEVHPRPGQDG AQ	

靶点信息

研究背景	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand fam ily. This protein is a ligand for the FN14/TWEAKR receptor. This cytokine has overlapping signaling functio ns with TNF, but displays a much wider tissue distribution. This cytokine, which exists in both membrane-bound and secreted forms, can induce apoptosis via multiple pathways of cell death in a cell type-specific manner. This cytokine is also found to promote proliferation and migration of endothelial cells, an d thus acts as a regulator of angiogenesis. Alternative splicing results in multiple transcript variants. Som e transcripts skip the last exon of this gene and continue into the second exon of the neighboring TNFSF1 3 gene; such read-through transcripts are contained in GeneID 407977, TNFSF12-TNFSF13.
基因 ID	8742
基因名	
Swiss	O43508
别名	TNFSF12;APO3L;DR3LG;TNLG4A;TWEAK

产品验证



Western blot analysis of TNFSF12 expressed in Rat kidney using TNFSF12 Rabbit pAb at 1:1000. Secondar y antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn