

FAM107A Rabbit pAb

货号: **B17678**

产品信息

反应	Human, Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	15kDa/17kDa/20kDa
实测分子量	18kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-144 of human FAM10 7A (NP_009108.1).
序列	MYSEIQRERADIGGLMARPEYREWNPELIKPKKLLNPVKASRSHQELHRELLMNHRRGLGVDSKPELQRVLEHRRRNQLIK KKKEELEAKRLQCPFEQELLRRQQRLNQLEKPPEKEEDHAPEFIKVRENLRRIATLTSEEREL

靶点信息

研究背景	Stress-inducible actin-binding protein that plays a role in synaptic and cognitive functions by modulating actin filamentous (F-actin dynamics. Mediates polymerization of globular actin to F-actin. Also binds to, st abilizes and bundles F-actin. Involved in synaptic function by regulating neurite outgrowth in an actin-dep endent manner and for the acquisition of hippocampus-dependent cognitive function, such as learning an d long-term memory (By similarity. Plays a role in the actin and microtubule cytoskeleton organization; n egatively regulates focal adhesion (FA assembly promoting malignant glial cell migration in an actin-, mic rotubule- and MAP1A-dependent manner. Also involved in neuroblastoma G1/S phase cell cycle progressi on and cell proliferation inhibition by stimulating ubiquitination of NF-kappa-B subunit RELA and NF-kappa -B degradation in a COMMD1- and actin-dependent manner. May play a role in tumor development.
基因 ID	11170
基因名	FAM107A
Swiss	O95990
别名	FAM107A;DRR1;TU3A

产品验证



Western blot analysis of FAM107A expressed in K-562 using FAM107A Rabbit pAb at 1:1000. Secondary an tibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% n on-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn